

# Superluminal Quantum Models of the Photon and Electron

<http://superluminalquantum.org>

Richard Gauthier, Santa Rosa, CA  
[rgauthier@qwickconnect.net](mailto:rgauthier@qwickconnect.net)

American Physical Society/ Regional Meeting  
Sacramento, California  
October 21-22, 2005


# **A single superluminal quantum composes a photon or an electron.**

A photon and an electron may be modeled by a point-like superluminal quantum moving in an open or closed helical trajectory, respectively, and characterized by an energy  $E$  and a forward momentum  $p$ , with a frequency  $\omega$  and a wavelength  $\lambda$  along its trajectory. Each superluminal quantum is associated with the usual quantum wave function which can produce interference and diffraction effects. A superluminal quantum is also point-like when it interacts with another superluminal quantum.

# The photon model is a superluminal uncharged quantum moving in an open helical trajectory.

A photon is modeled as a helical movement of a superluminal quantum with forward momentum  $p$  along a helical path of radius  $R$ , pitch (wavelength)  $\lambda$ , and forward helical angle  $\theta$ . By combining the angular momentum and linear momentum relations for a photon with this spatially extended helical geometry, the result is:

- 1) The forward helical angle  $\theta$  of the superluminal quantum is found to be 45 degrees, for any photon wavelength.
- 2) The superluminal quantum's helical radius is found to always be  $R = \lambda / 2\pi$ .
- 3) The speed of the superluminal quantum is  $\sqrt{2}c = 1.414..c$  along the helical path.



A photon is modeled by an uncharged superluminal quantum moving at  $1.414c$  along an open 45-degree helical trajectory of radius  $R = \lambda / 2\pi$ .

# Proof of the above three statements for the superluminal quantum model of the photon.

Assume all the momentum  $P$  of the superluminal quantum along its helical path is concentrated at a single point on the helical path. This instantaneous momentum  $P$  at this point is directed along its helical path, which makes a constant forward angle  $\theta$  with the longitudinal direction of the photon. So the longitudinal component of the superluminal quantum's momentum is  $P \cos(\theta) = 2\pi\hbar / \lambda$ , the experimental longitudinal or linear momentum of a photon. The momentum  $P$ 's transverse component (which is also perpendicular to the helical radius  $R$  to this point object from the helical axis) of momentum is  $P \sin(\theta)$ , so the angular momentum or spin of this simple photon model is  $S = RP \sin(\theta) = \hbar$ , the experimental spin or angular momentum of the photon. Combining these two equations containing  $\theta$  gives  $\sin(\theta) / \cos(\theta) = \tan(\theta) = \lambda / 2\pi R$  (the values  $\hbar$  and  $P$  cancel out). Now consider the helical geometry. As the superluminal quantum advances a distance  $\lambda$  in the longitudinal direction, it moves a transverse distance  $2\pi R$ , i.e. once around the circle of radius  $R$  defined by the helix. From the way  $\theta$  is defined,  $\tan(\theta)$  equals this transverse distance divided by the longitudinal distance traveled in one wavelength, or  $\tan(\theta) = 2\pi R / \lambda$ . So we now have two equations for  $\tan(\theta)$ . So  $\tan(\theta) = 2\pi R / \lambda = \lambda / 2\pi R$ . This will only be true if  $\lambda = 2\pi R$ , that is, when  $R = \lambda / 2\pi$  and  $\theta = 45^\circ$ . So since  $\theta = 45^\circ$  and the forward velocity of the photon is  $c$ , the speed of the superluminal quantum is  $c / \cos(45^\circ) = c / .707 = 1.414c$ .


# The electron model is a superluminal charged quantum moving in a closed double-looped helical trajectory.

The electron is modeled by a closed double-looped helical trajectory of a superluminal quantum of charge  $-e$  and energy  $E = mc^2$ . Consider one turn of a helix of pitch one Compton wavelength to be looped twice before joining its ends together. It will have a double-looped circular axis of total length one Compton wavelength. If the radius of its circular axis is  $R_0 = \frac{1}{2} \hbar / mc$  and the radius of the closed helix is  $R_0 \sqrt{2}$ , then the equation for the coordinates of the superluminal quantum as it moves around this closed helix with angular frequency  $\omega_0 = mc^2 / \hbar$  is

$$x(t) = R_0 (1 + \sqrt{2} \cos(\omega_0 t)) \cos(2\omega_0 t)$$

$$y(t) = R_0 (1 + \sqrt{2} \cos(\omega_0 t)) \sin(2\omega_0 t)$$

$$z(t) = R_0 \sqrt{2} \sin(\omega_0 t)$$



**Two views of the superluminal quantum model of the electron. The circle is the axis of the double-looped helical trajectory of the superluminal quantum.**

# Properties the superluminal quantum model of the electron shares with the Dirac equation's electron with *Zitterbewegung*

1. The calculated electron spin  $s = \frac{1}{2} \hbar$ .
2. The calculated electron magnetic moment  $\mu = e\hbar / 2mc$  the Bohr magneton ( $g=2$ ).
3. The *Zitterbewegung* light velocity  $c$  of the electron (the electron model resembles a circling photon model with speed  $c$ ).
4. The *Zitterbewegung* frequency  $2\omega_0$ , where  $\omega_0 = mc^2 / \hbar$ .
5. The *Zitterbewegung* radius  $R_0 = \frac{1}{2} \hbar / mc$ .
6. The prediction of the electron and the positron (due to the two possible helicities in the electron model).
7. The *Zitterbewegung* distinction between the coordinates of the electron's point charge and the position of the electron as a whole.
8. The non-conservation of momentum of the *Zitterbewegung* motion.